



TITLE:

Role of clock genes in insulin secretion

AUTHOR(S):

Harada, Norio; Inagaki, Nobuya

CITATION:

Harada, Norio ...[et al]. Role of clock genes in insulin secretion. Journal of Diabetes Investigation 2016, 7(6): 822-823

ISSUE DATE:

2016-11

URL:

<http://hdl.handle.net/2433/217349>

RIGHT:

© 2016 The Authors. Journal of Diabetes Investigation published by Asian Association for the Study of Diabetes (AASD) and John Wiley & Sons Australia, Ltd This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Role of clock genes in insulin secretion

Circadian locomotor output cycles kaput (CLOCK) and brain and muscle aryl hydrocarbon receptor nuclear translocator-like protein 1 (BMAL1) are master clock genes that regulate circadian rhythm in the hypothalamus and peripheral tissues in mammals; and control not only sleep cycle, but also many other physiological functions, such as body temperature, heart rate and hormone secretion. CLOCK and BMAL1 protein form a heterodimer and bind to enhancer box (E-box) elements located upstream of circadian rhythm-related genes, which are period (PER) and cryptochrome (CRY), and non-circadian rhythm-related genes, resulting in production of PER, CRY and other non-circadian rhythm-related proteins. In the cytoplasm, PER and CRY protein form a heterodimer that is subsequently translocated into the nucleus and inhibits CLOCK and BMAL1-induced transcriptions. This negative feedback loop is an important part of mammalian circadian rhythm (Figure 1)¹.

Recent studies of CLOCK-mutant and BMAL1-knockout mice show that circadian rhythm influences the development of metabolic syndrome. Locomotor activity of CLOCK-mutant mice was higher than that of wild-type mice in the light phase condition, and the feeding pattern of the mutant mice was apparently different from that of wild-type mice². Energy expenditure was decreased and bodyweight was increased in the mutant mice compared with that in wild-type mice. Furthermore, plasma triglyceride and low-density lipoprotein cholesterol concentrations were decreased in mutant mice. In contrast, bodyweight and adipose tissue size were significantly decreased in systemic BMAL1-knockout

mice compared with wild-type mice.³ BMAL1-knockout mice had higher plasma triglyceride and low-density lipoprotein cholesterol concentrations compared with wild-type mice. These results show that CLOCK and BMAL1 are involved in lipid metabolism and bodyweight control. However, there was a large difference in phenotype between CLOCK-mutant and BMAL1-knockout mice. It is speculated that CLOCK and BMAL1 regulates different non-circadian proteins, which are associated with lipid metabolism and obesity.

Previously, β -cell-specific BMAL1-knockout mice were generated to evaluate the effect of BMAL1 on insulin secretion⁴. Plasma insulin concentrations after intraperitoneal glucose injection were significantly lower in β -cell-specific BMAL1-knockout mice compared with those in wild-type mice, resulting in hyperglycemia. Insulin secretion in response to glucose, adenylyl cyclase activator (forskolin), glucagon-like peptide-1 receptor agonist (exendin-4), cyclic adenosine 3', 5'-monophosphate (8-bromo-cyclic adenosine 3', 5'-monophosphate)

and hyperdepolarization (KCl) were significantly decreased in the isolated islets of β -cell-specific BMAL1-knockout mice compared with wild-type mice. These results show that BMAL1 is also involved in insulin secretion. However, the detailed mechanism of BMAL1-mediated regulation of insulin secretion from pancreatic β -cells is unclear. Perelis *et al.*⁵ clearly showed the regulatory role of BMAL1 in insulin secretion by genome-wide analysis of isolated islets. They generated tamoxifen-induced β -cell-specific BMAL1-knockout mice, and showed that acquired BMAL1 deficiency in β -cells decreases insulin secretion in response to glucose, forskolin, 8-bromo-cyclic adenosine 3', 5'-monophosphate and KCl. Genomic-wide analysis using ribonucleic acid sequence suggested reduction of not only circadian rhythm-related genes, but also genes associated with transport and membrane fusion of insulin vesicle in the islets. Additionally, data of chromatin immunoprecipitation-sequence showed that CLOCK and BMAL1 bind at distal regulatory sites of circadian rhythm-related genes in β -cells. This region also




Figure 1 | Molecular mechanism of circadian rhythm. Circadian locomotor output cycles kaput (CLOCK) and brain and muscle aryl hydrocarbon receptor nuclear translocator-like protein 1 (BMAL1) protein form a heterodimer and bind to E-box elements, resulting in induction of period (PER), cryptochrome (CRY) and non-circadian rhythm-related gene transcriptions. In the cytoplasm, PER and CRY protein is subsequently translocated into the nucleus, and inhibits CLOCK and BMAL1-induced transcriptions (negative feedback). E-box, enhancer box.

*Corresponding author, Norio Harada

Tel: +81-75-751-3560

Fax: +81-75-751-4244

E-mail address: nharada@kuhp.kyoto-u.ac.jp

Received 2 March 2016; revised 7 March 2016;

accepted 8 March 2016

contained the active enhancer at which pancreatic transcriptional factor pancreatic and duodenal homeobox 1 bind. Thus, clock genes regulate the genes (Pdx1) that are associated with insulin secretion and production in β -cells.

DISCLOSURE

The authors declare no conflict of interest.

Norio Harada*, Nobuya Inagaki
*Diabetes, Endocrinology and Nutrition,
Graduate School of Medicine, Kyoto
University, Kyoto, Japan*

REFERENCES

1. Preitner N, Damiola F, Lopez-Molina L, *et al.* The orphan nuclear receptor REV-ERB α controls circadian transcription within the positive limb of the mammalian circadian oscillator. *Cell* 2002; 110: 251–260.
2. Turek FW, Joshu C, Kohsaka A, *et al.* Obesity and metabolic syndrome in circadian Clock mutant mice. *Science* 2005; 308: 1043–1045.
3. Shimba S, Ogawa T, Hitosugi S, *et al.* Deficient of a clock gene, brain and muscle Arnt-like protein-1 (BMAL1), induces dyslipidemia and ectopic fat formation. *PLoS One* 2011; 6: e25231.
4. Marcheva B, Ramsey KM, Buhr ED, *et al.* Disruption of the clock components CLOCK and BMAL1 leads to hypoinsulinaemia and diabetes. *Nature* 2010; 466: 627–631.
5. Perelis M, Marcheva B, Ramsey KM, *et al.* Pancreatic β cell enhancers regulate rhythmic transcription of genes controlling insulin secretion. *Science* 2015; 350: aac4250.

Doi: 10.1111/jdi.12517